Acta Mathematica Sinica, English Series Mar., 2017, Vol. 33, No. 3, pp. 341–361 Published online: July 20, 2016 DOI: 10.1007/s10114-016-6029-7 Http://www.ActaMath.com

Acta Mathematica Sinica, English Series

Springer-Verlag Berlin Heidelberg & The Editorial Office of AMS 2017

Representations of Frobenius-type Triangular Matrix Algebras

Fang LI Chang $YE^{(1)}$

Department of Mathematics, Zhejiang University, Hangzhou 310027, P. R. China E-mail: fangli@zju.edu.cn yc2009@zju.edu.cn

Abstract The aim of this paper is mainly to build a new representation-theoretic realization of finite root systems through the so-called Frobenius-type triangular matrix algebras by the method of reflection functors over any field. Finally, we give an analog of APR-tilting module for this class of algebras. The major conclusions contains the known results as special cases, e.g., that for path algebras over an algebraically closed field and for path algebras with relations from symmetrizable cartan matrices. Meanwhile, it means the corresponding results for some other important classes of algebras, that is, the path algebras of quivers over Frobenius algebras and the generalized path algebras endowed by Frobenius algebras at vertices.

Keywords Frobenius-type triangular matrix algebras, reflection functor, locally free module, root system, APR-tilting module

MR(2010) Subject Classification 16G10, 16G20, 16G60

1 Introduction and Preliminaries

Let Q be a finite connected acyclic quiver, and let H = kQ be the path algebra of Q for an algebraically closed field k. Gabriel showed that the quiver Q is representation-finite if and only if Q is a Dynkin quiver of type A_n, D_n, E_6, E_7, E_8 in [8]. In this case, there is a bijection between the isomorphism classes of indecomposable representations of Q and the set of positive roots of the corresponding simple complex Lie algebra. Bernštein et al. introduced the machinery of Coxeter functors, which are defined as compositions of reflection functors, to give an elegant proof of Gabriel's theorem in [5]. Gabriel also showed that there are functorial isomorphisms $SC^{\pm}(-) \cong \tau^{\pm}(-)$ in [9], where S is a twist functor, C^{\pm} are the Coxeter functors and $\tau(-)$ is the Auslander–Reiten translation. Auslander et al. showed that there exists an H-module T satisfying the functorial isomorphisms $F_k^{\pm}(-) \cong \operatorname{Hom}_H(T,-)$ in [4] for the BGP-reflection functors F_k^{\pm} and the APR-tilting module T.

Some of these results have been developed to valued graphs or k-species by Dlab and Ringel for a field k, see [6, 7, 18]. Moreover, in [10], for any field k, Geiss et al. generalized them to a class of 1-Gorenstein algebras A, which were defined via quivers with relations associated to symmetrizable Cartan matrices, as follows.

Supported by National Natural Science Foundation of China (Grant Nos. 11271318 and 11571173); and the Zhejiang Provincial Natural Science Foundation of China (Grant No. LZ13A010001)

1) Corresponding author

Received January 12, 2016, accepted May 11, 2016

Let $C = (c_{ij}) \in M_n(\mathbb{Z})$ be a symmetrizable generalized Cartan matrix with a symmetrizer $D = \text{diag}(c_1, \ldots, c_n)$. For all $c_{ij} < 0$, write that $g_{ij} := |\text{gcd}(c_{ij}, c_{ji})|, f_{ij} := |c_{ij}|/g_{ij}, k_{ij} := \text{gcd}(c_i, c_j)$.

An orientation of C is a subset $\Omega \subset \{1, 2, ..., n\} \times \{1, 2, ..., n\}$ such that the following hold: (i) $\{(i, j), (j, i)\} \cap \Omega \neq \emptyset$ if and only if $c_{ij} < 0$;

(ii) For each sequence $((i_1, i_2), (i_2, i_3), \dots, (i_t, i_{t+1}))$ with $t \ge 1$ and $(i_s, i_{s+1}) \in \Omega$ for all $1 \le s \le t$ we have $i_1 \ne i_{t+1}$.

For an orientation Ω of C, let $Q := Q(C, \Omega) := (Q_0, Q_1, s, t)$ be the quiver with the set of vertices $Q_0 := \{1, \ldots, n\}$ and the set of arrows

$$Q_1 := \{\alpha_{ij}^{(g)} : j \to i | (i,j) \in \Omega, 1 \le g \le g_{ij}\} \cup \{\varepsilon_i : i \to i | 1 \le i \le n\}.$$

For a quiver $Q = Q(C, \Omega)$ and a symmetrizer $D = \text{diag}(c_1, \ldots, c_n)$ of C, let

$$A = A(C, D, \Omega) := kQ/I, \tag{1.1}$$

where kQ is the path algebra of Q, and I is the ideal of kQ defined by the following relations:

(i) For each *i*, we have the nilpotency relation $\varepsilon_i^{c_i} = 0$.

(ii) For each $(i, j) \in \Omega$ and each $1 \leq g \leq g_{ij}$, we have the commutativity relation $\varepsilon_i^{f_{ji}} \alpha_{ij}^{(g)} = \alpha_{ij}^{(g)} \varepsilon_j^{f_{ij}}$.

It was proved in [10] that the algebra A, given in (1.1) is 1-Gorenstein.

The aim of this paper is to give a larger class of 1-Gorenstein algebras in which the important conclusions about representation theory still hold. More specifically, using of the dual basis lemma and the method of reflection functors which developed by Gabriel et al., we will build a new representation-theoretic realization of finite root systems through Frobenius-type triangular matrix algebras.

In the sequel, the ground field k is always permitted to be any field.

For $n \geq 2$, define a triangular matrix algebra Γ of order n satisfying

$$\Gamma = \begin{pmatrix} A_1 & A_{12} & \dots & A_{1n} \\ 0 & A_2 & \dots & A_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & A_n \end{pmatrix},$$

where each A_i is an algebra and A_{ij} an $A_i \cdot A_j$ -bimodule with bimodule maps $\mu_{ilj} : A_{il} \otimes_{A_l} A_{lj} \rightarrow A_{ij}$ such that the following diagram commutes:

$$\begin{array}{c|c} A_{il} \otimes_{A_l} A_{lj} \otimes_{A_j} A_{jt} & \xrightarrow{\mu_{ilj} \otimes id_{A_{jt}}} & A_{ij} \otimes_{A_j} A_{jt} \\ id_{A_{il}} \otimes_{\mu_{ljt}} \downarrow & & \downarrow \\ A_{il} \otimes_{A_l} A_{lt} & \xrightarrow{\mu_{ilt}} & A_{it} \end{array}$$

for $1 \leq i < l < j < t \leq n$, whose multiplication is given by $(AB)_{ij} = \sum_{i < l < j} \mu_{ilj}(a_{il} \otimes b_{lj})$ for $A = (a_{ij})_{n \times n}, B = (b_{ij})_{n \times n} \in \Gamma$, where $(AB)_{ij}$ means the (i, j)-entry of AB.

A representation X of Γ is defined as a datum

$$X = \begin{pmatrix} X_1 \\ X_2 \\ \vdots \\ X_n \end{pmatrix}_{\phi_{ij}}$$

,

with ϕ_{ij} : $A_{ij} \otimes_{A_j} X_j \to X_i$ an A_i -module morphism for $1 \le i < j \le n$ so that it satisfies the following commutative diagram:

~ · 1

A morphism f from a representation X to another representation Y is defined as a datum $(f_i, 1 \leq i \leq n)$, where $f_i : X_i \to Y_i$ is an A_i -map such that $f_i \phi_{ij} = \phi_{ij} (id_{A_{i,j}} \otimes f_j)$ for each $1 \leq i < j \leq n$.

Then we obtain the representation category of Γ , denoted as $\operatorname{Rep}(\Gamma)$.

Recall that a finite dimensional algebra A is said to be a Frobenius algebra if there exists an isomorphism of left modules ${}_{A}A \cong D(A_{A})$, where $D := \operatorname{Hom}_{k}(-, k)$. See [17].

Definition 1.1 For each i = 1, ..., n, let A_i be a Frobenius algebra with unit e_i . For each $1 \leq i < j \leq n$, let B_{ij} be an A_i - A_j -bimodule such that B_{ij} is a free left A_i -module and free right A_j -module of finite rank respectively. Suppose there are A_i - A_j -bimodule isomorphisms $\operatorname{Hom}_{A_i}(B_{ij}, A_i) \cong \operatorname{Hom}_{A_j}(B_{ij}, A_j)$ for all $1 \leq i < j \leq n$. Let

$$A_{ij} = \bigoplus_{l=0}^{j-i-1} \bigoplus_{i < k_1 < k_2 < \dots < k_l < j} B_{ik_1} \otimes_{A_{k_1}} B_{k_1 k_2} \otimes \dots \otimes_{A_{k_l}} B_{k_l j}$$
(1.2)

for $1 \leq i < j \leq n$, where l = 0 means the direct summand B_{ij} .

Define a triangular matrix algebra

$$\Lambda = \begin{pmatrix} A_1 & A_{12} & \dots & A_{1n} \\ 0 & A_2 & \dots & A_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & A_n \end{pmatrix}$$

which is called a Frobenius-type triangular matrix algebra, if it satisfies (1.2) with bimodule maps $\mu_{ijq}: A_{ij} \otimes_{A_j} A_{jq} \to A_{iq}$ are the natural inclusion maps.

The algebra $A = A(C, D, \Omega) = kQ/I$ in (1.1) from [10] is indeed a Frobenius-type triangular matrix algebra. To see this, one needs only to re-order vertexes by i < j if $(i, j) \in \Omega$ and take $A_i = e_i A e_i, \ B_{ij} = A_i \text{Span}_k(\alpha_{ij}^{(g)}|1 \le g \le g_{ij})A_j$. It is easy to see that such A satisfies the condition of Frobenius-type triangular matrix algebra in Definition 1.1.

Following this fact, in this paper the major results on Frobenius-type triangular matrix algebras are the improvement of the corresponding ones in [10]. For convenience, in the sequel, we will always assume A_i to be finite dimensional for all i = 1, ..., n.

Remark 1.2 (i) In Definition 1.1, if each A_i is only a finite dimensional algebra, and the condition $\operatorname{Hom}_{A_i}(B_{ij}, A_i) \cong \operatorname{Hom}_{A_j}(B_{ij}, A_j)$ is replaced by that $\operatorname{Hom}_{A_i}(B_{ij}, A_i)$ is a projective A_j -module or an injective A_j -module, then Λ is called a *normally upper triangular gm algebra*, which was introduced and investigated in [15].

Since $\operatorname{Hom}_{A_j}(B_{ij}, A_j)$ is a free left A_j -module and $\operatorname{Hom}_{A_i}(B_{ij}, A_i) \cong \operatorname{Hom}_{A_j}(B_{ij}, A_j)$, we obtain that $\operatorname{Hom}_{A_i}(B_{ij}, A_i)$ is a free left A_j -module, then, of course, a projective A_j -module. So all Frobenius-type triangular matrix algebras are normally upper triangular gm algebras.

(ii) In Definition 1.1, let $B_{ij} = 0$ for all $|i - j| \ge 2$, and let A_i be any rings. Then we obtain a class of triangular matrix rings which were studied in [20] about their module categories and some homological characterizations.

(iii) A path algebra of quiver over algebra $\Lambda = AQ = A \bigotimes_k kQ$ for an acyclic quiver Q and a Frobenius algebra A studied in [19] is also a Frobenius-type triangular matrix algebra. Here one needs only to take all $A_i = A$ and $B_{ij} = \bigoplus_{s=1}^{\#\{\alpha:j\to i\}} A$. In particular, it was investigated in [19] for the case $A = k[T]/[T^2]$ the algebra of dual numbers.

(iv) A generalized path algebra $\Lambda = k(Q, \mathcal{A})$, with an acyclic quiver Q, $\mathcal{A} = \{A_i\}_{i \in Q_0}$ and Forbenius algebras A_i ($i \in Q_0$), is a Frobenius-type triangular matrix algebra through taking all B_{ij} to be generalized arrows from j to i and thus obtaining A_{ij} as generalized paths from jto i. For more details, see [15].

Let

$$X = \begin{pmatrix} X_1 \\ X_2 \\ \vdots \\ X_n \end{pmatrix}_{\phi}$$

be a Λ -module. It is easy to see that $\phi_{ij}: A_{ij} \bigotimes_{A_j} X_j \to X_i$ is uniquely determined by $\phi_{ij}|_{\operatorname{Res}(B_{ij} \bigotimes X_j)}: B_{ij} \bigotimes_{A_j} X_j \to X_i$ for $1 \leq i < j \leq n$.

Remark 1.3 Without ambiguity, we sometimes omit to write ϕ_{ij} , especially when ϕ_{ij} is a natural inclusion for all $1 \le i < j \le n$.

Remark 1.4 In this paper, we consider only Λ which is connected, that is, Λ can NOT be written as

$$\left(\begin{array}{cc} \Lambda_1 & 0\\ 0 & \Lambda_2 \end{array}\right)$$

for two non-zero Frobenius-type triangular matrix algebras Λ_1 and Λ_2 . Otherwise, for

$$\Lambda = \left(\begin{array}{cc} \Lambda_1 & 0\\ 0 & \Lambda_2 \end{array} \right),$$

any representation M of Λ can be obtained from representations M_i of Λ_i (i = 1, 2). Roughly speaking,

$$M = \left(\begin{array}{c} M_1 \\ 0 \end{array}\right) \oplus \left(\begin{array}{c} 0 \\ M_2 \end{array}\right).$$

Our main result is as follows.

Theorem 1.5 For a Frobenius-type triangular matrix algebra Λ , the following statements hold:

(a) The number of isomorphism classes of indecomposable τ -locally free Λ -modules is finite if and only if C is of Dynkin type.

(b) If C is of Dynkin type, then the mapping $\underline{\operatorname{rank}} : X \mapsto \underline{\operatorname{rank}}(X)$ induces a bijection between the set of isomorphism classes of indecomposable τ -locally free Λ -modules and the set of positive roots of the quadratic form $q_C(x)$.

Theorem 1.6 For a Frobenius-type triangular matrix algebra Λ , there is a functorial isomorphism

$$F_1^+(-) \cong \operatorname{Hom}_{\Lambda}(T_1, -) : \operatorname{rep}(\Lambda) \to \operatorname{rep}(S_1(\Lambda)).$$

This article is organized as follows. In Section 2, we study some properties of Frobeniustype triangular matrix algebras and in particular, show that they are a class of 1-Gorenstein algebras. In Section 3, we define the reflection functors for Frobenius-type triangular matrix algebras and give the relation between Coxeter functors and Auslander–Reiten translation for these algebras. In Section 4, we recall some definitions and basic facts on Cartan matrices, quadratic forms and Weyl groups. Then we give a new representation-theoretic realizations of all finite root systems via Frobenius-type triangular matrix algebras. In Section 5, we study the generalized versions of APR-tilting modules over Frobenius-type triangular matrix algebras.

One of our motivation is, in the further work, using the results in this paper, to characterise the categorification of cluster algebras in the case for skew-symmetrizable type.

2 Locally Free Modules and 1-Gorenstein Property

Proposition 2.1 ([15]) The representation category of a Frobenius-type triangular matrix algebra Λ and the module category of Λ are equivalent.

We denote $B_{ji} = \operatorname{Hom}_{A_i}(B_{ij}, A_i) \cong \operatorname{Hom}_{A_j}(B_{ij}, A_j)$ for $1 \le i < j \le n$.

Proposition 2.2 For $1 \le i < j \le n$, B_{ji} is a free left A_j -module and a free right A_i -module. Also, we have $B_{ij} = \operatorname{Hom}_{A_i}(B_{ji}, A_i) \cong \operatorname{Hom}_{A_j}(B_{ji}, A_j)$.

Proof By the Frobeniusness of A_i , we have the isomorphism $\operatorname{Hom}_{A_i}(\operatorname{Hom}_{A_i}(-, A_i), A_i) \cong 1$. Then the result follows.

Proposition 2.3 For any A_j -module M_j and A_i -module N_i , we have the isomorphism

$$\operatorname{Hom}_{A_i}\left(B_{ij}\bigotimes_{A_j}M_j,N_i\right)\cong\operatorname{Hom}_{A_j}\left(M_j,B_{ji}\bigotimes_{A_i}N_i\right).$$

Proof The adjunction map gives an isomorphism of k-vector spaces:

$$\operatorname{Hom}_{A_i}\left(B_{ij}\bigotimes_{A_j}M_j,N_i\right)\cong\operatorname{Hom}_{A_j}(M_j,\operatorname{Hom}_{A_i}(B_{ij},N_i)).$$

Since $\operatorname{Hom}_{A_i}(B_{ij}, N_i) \cong \operatorname{Hom}_{A_i}(B_{ij}, A_i \bigotimes_{A_i} N_i) \stackrel{(*)}{\cong} \operatorname{Hom}_{A_i}(B_{ij}, A_i) \bigotimes_{A_i} N_i \cong B_{ji} \bigotimes_{A_i} N_i$, we have

$$\operatorname{Hom}_{A_i}\left(B_{ij}\bigotimes_{A_j}M_j,N_i\right)\cong\operatorname{Hom}_{A_j}\left(M_j,B_{ji}\bigotimes_{A_i}N_i\right),$$

where the second isomorphism (*) follows from that B_{ij} is a finitely generated free A_i -module. **Lemma 2.4** (The Dual Basis Lemma) Let A be an algebra, and P be a finitely generated projective A-module, $\operatorname{Hom}_A(P, A)$. Then there exist $x_1, \ldots, x_m \in P, f_1, \ldots, f_m \in \operatorname{Hom}_A(P, A)$, such that for each $x \in P, x = \sum_{i=1}^m f_i(x)x_i$.

Since B_{ij} is a finitely generated projective left A_i -module and a finitely generated projective right A_j -module, by the above lemma, there exist $L_{ij} \subseteq B_{ij}$, $L_{ij}^* \subseteq \text{Hom}_{A_i}(B_{ij}, A_i)$ and $R_{ij} \subseteq B_{ij}$, $R_{ij}^* \subseteq \text{Hom}_{A_j}(B_{ij}, A_j)$ such that for each $b_{ij} \in B_{ij}$, $b_{ij} = \sum_{\ell \in L_{ij}} \ell^*(b_{ij})\ell = \sum_{r \in R_{ij}} r^*(b_{ij})r$.

For the isomorphism in Proposition 2.3, for each $\phi_{ij} \in \operatorname{Hom}_{A_i}(B_{ij} \bigotimes_{A_j} M_j, N_i)$, we have $\overline{\phi_{ij}} \in \operatorname{Hom}_{A_j}(M_j, B_{ji} \bigotimes_{A_i} N_i)$ satisfying $\overline{\phi_{ij}}(m_j) = \sum_{\ell \in L_{ij}} \ell^* \otimes \phi_{ij}(\ell \otimes m_j)$.

On the other hand, for each $\psi_{ij} \in \operatorname{Hom}_{A_j}(M_j, B_{ji} \bigotimes_{A_i} N_i)$, we have $\overline{\psi_{ij}} \in \operatorname{Hom}_{A_i}(B_{ij} \bigotimes_{A_j} M_j, N_i)$ satisfying $\overline{\psi_{ij}}(b_{ij} \otimes m_j) = \sum_{\ell \in L_{ij}} \ell^*(b_{ij})\psi_{ij}(m_j)_{\ell}$, where the elements $\psi_{ij}(m_j)_l \in N_i$ are uniquely determined by $\psi_{ij}(m_j) = \sum_{\ell \in L_{ij}} \ell^* \otimes \psi_{ij}(m_j)_{\ell}$.

Let Λ be a Frobenius-type triangular matrix algebra as in Definition 1.1. Denote $P_i = \Lambda e_i$ the projective Λ -module for the idempotent e_i as the unit 1 of A_i and I_i the corresponding injective Λ -module for $1 \leq i \leq n$. Obviously,

$$P_i = (A_{1i}, A_{2i}, \dots, A_{ii}, 0, \dots, 0)^t.$$
(2.1)

Also, we denote E_i the Λ -module $(0, \ldots, A_i, \ldots, 0)_{\phi_{lj}}^t$, where A_i is in the *i*-th row and all $\phi_{lj} = 0$.

Lemma 2.5 For two k-algebras A and B, assume that M is an A-B-bimodule such that M is a projective left A-module, and P is a projective left B-module. Then $M \bigotimes_B P$ is a projective left A-module.

Proposition 2.6 For $E_i = (0, \ldots, A_i, \ldots, 0)_{\phi_{lj}}^t$ with $i = 1, \ldots, n$, it holds that $\operatorname{proj.dim}(E_i) \leq 1$ and $\operatorname{inj.dim}(E_i) \leq 1$.

Proof Clearly, $E_1 = P_1$. For every i = 2, ..., n, we have exact sequences:

$$0 \to \bigoplus_{j=1}^{i-1} P_j \bigotimes_{A_j} B_{ji} \to P_i \to E_i \to 0.$$
(2.2)

So, $\operatorname{proj.dim}(E_i) \leq 1$ by Lemma 2.5. Let E'_i be the right Λ -module such that $D(E'_i) \cong E_i$. For $i = 1, \ldots, n-1$, since A_i is a Frobenius algebra, there is a canonical exact sequence

$$0 \to \bigoplus_{j=i+1}^{n} B_{ij} \bigotimes_{A_j} e_j \Lambda \to e_i \Lambda \to E'_i \to 0.$$
(2.3)

Applying the duality D to (2.3), we get a minimal injective resolution

$$0 \to E_i \to D(e_i\Lambda) \to \bigoplus_{j=i+1}^n D\left(B_{ij}\bigotimes_{A_j} e_j\Lambda\right) \to 0.$$
(2.4)

Clearly, E_n is injective. So, $\operatorname{inj.dim}(E_i) \leq 1$ for $1 \leq i \leq n$.

Definition 2.7 For a Frobenius-type triangular matrix algebra Λ , following [10], a finitely generated Λ -representation X is called locally free if X_i are free A_i -modules for all $1 \leq i \leq n$.

Denote by $\operatorname{rep}_{l.f.}(\Lambda)$ the subcategory of all locally free Λ -modules.

Corollary 2.8 For a Frobenius-type triangular matrix algebra Λ , suppose $X \in \operatorname{rep}_{l.f.}(\Lambda)$, then it holds that $\operatorname{proj.dim}(X) \leq 1$ and $\operatorname{inj.dim}(X) \leq 1$.

Proof We have a short exact sequence

$$0 \to e_1 X \to X \to (1 - e_1) X \to 0,$$

where e_1X and $(1 - e_1)X$ are locally free. By Proposition 2.6 and using induction on n, we know that the projective and the injective dimensions of e_1X and $(1 - e_1)X$ are at most one. **Remark 2.9** For the algebra $A = A(C, D, \Omega)$ in [10], the three conditions

 $X \in \operatorname{rep}_{l,f}(A), \quad \operatorname{proj.dim}(X) \le 1, \quad \operatorname{inj.dim}(X) \le 1$

are equivalent. But it is generally not true for a Frobenius-type triangular matrix algebra.

An algebra A is called *m*-Gorenstein if $inj.dim(A) \leq m$ and $proj.dim(DA) \leq m$. Such algebras were firstly introduced and studied in [11].

Corollary 2.10 The Frobenius-type triangular matrix algebra Λ is a 1-Gorenstein algebra.

Proof It is a direct consequence of Corollary 2.8.

More general discussion related to this corollary can also be found in [21], in the various way.

Recall that for an algebra A, an A-module X is τ -rigid (resp. τ^- -rigid) if $\operatorname{Hom}_A(X, \tau(X)) = 0$ (resp. $\operatorname{Hom}_A(\tau^-(X), X) = 0$). We call the Λ -mod X rigid if $\operatorname{Ext}^1_{\Lambda}(X, X) = 0$. See [1].

When proj.dim $(X) \leq 1$, we have a functorial isomorphism $\operatorname{Ext}_{A}^{1}(X,Y) \cong D\operatorname{Hom}_{A}(Y,\tau(X));$ when inj.dim $(X) \leq 1$, we have a functorial isomorphism $\operatorname{Ext}_{A}^{1}(X,Y) \cong D\operatorname{Hom}_{A}(\tau^{-}(Y),X).$ See [3].

Corollary 2.11 For a Frobenius-type triangular matrix algebra Λ , let $X \in \operatorname{rep}_{l.f.}(\Lambda)$. Then X is rigid if and only if X is τ -rigid, also if and only if X is τ -rigid.

Proposition 2.12 For a Frobenius-type triangular matrix algebra Λ , the subcategory $\operatorname{rep}_{l.f.}(\Lambda)$ is closed under extensions, kernels of epimorphisms and cokernels of monomorphisms.

Proof Let $0 \to X \xrightarrow{f} Y \xrightarrow{g} Z \to 0$ be a short exact sequence in rep(Λ). For each $1 \leq i \leq n$, this induces a short exact sequence

$$0 \to e_i X \xrightarrow{f} e_i Y \xrightarrow{g} e_i Z \to 0 \tag{2.5}$$

of left A_i -modules.

By the definition, when $X, Z \in \operatorname{rep}_{l.f.}(\Lambda)$, $e_i X, e_i Z$ are both A_i -free, and then $e_i X$ is injective via A_i is a Frobenius algebra and $e_i Z$ is projective for any *i*. Hence, the short exact sequence (2.5) splits, i.e., $e_i Y \cong e_i X \bigoplus e_i Z$. It follows that each $e_i Y$ is free A_i -module and then $Y \in \operatorname{rep}_{l.f.}(\Lambda)$.

When $Y, Z \in \operatorname{rep}_{l.f.}(\Lambda)$, $e_i Y, e_i Z$ are both A_i -free and then $e_i Z$ is a projective module. Hence (2.5) splits, i.e., $e_i Y \cong e_i X \bigoplus e_i Z$. By the Krull–Schmidt theorem, it is easy to see that each $e_i X$ is a free A_i -module. So, $X \in \operatorname{rep}_{l.f.}(\Lambda)$.

When $X, Y \in \operatorname{rep}_{l.f.}(\Lambda)$, $e_i X, e_i Y$ are both A_i -free and then $e_i X$ is injective via A_i is a Frobenius algebra. Hence (2.5) splits. Similarly, by the Krull–Schmidt theorem, each $e_i Z$ is free A_i -module. So, $Z \in \operatorname{rep}_{l.f.}(\Lambda)$.

3 Reflection Functors and AR-translation

Let

$$\Lambda = \begin{pmatrix} A_1 & A_{12} & \dots & A_{1n} \\ 0 & A_2 & \dots & A_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & A_n \end{pmatrix}$$

be a Frobenius-type triangular matrix algebra. Denote

$$S_1(\Lambda) = \begin{pmatrix} A_2 & A_{23} & \dots & A_{2n} & A_{21} \\ 0 & A_3 & \dots & A_{3n} & A_{31} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & A_n & A_{n1} \\ 0 & 0 & \dots & 0 & A_1 \end{pmatrix},$$

where $B_{j1} = \operatorname{Hom}_{A_1}(B_{1j}, A_1)$ and

$$A_{j1} = \bigoplus_{l=0}^{n-j} \bigoplus_{j < k_1 < k_2 < \dots < k_l \le n} B_{jk_1} \bigotimes_{A_{k_1}} B_{k_1k_2} \bigotimes \dots \bigotimes_{A_{k_l}} B_{k_l 1}$$

for j = 2, ..., n.

 B_{j1} is a free right A_1 -module and also a free left A_j -module, since $B_{j1} = \operatorname{Hom}_{A_1}(B_{1j}, A_1) \cong \operatorname{Hom}_{A_j}(B_{1j}, A_j)$. For $S_1(\Lambda)$, we have

$$\operatorname{Hom}_{A_1}(B_{j1}, A_1) \cong \operatorname{Hom}_{A_1}(\operatorname{Hom}_{A_1}(B_{1j}, A_1), A_1) \cong B_{1j}$$
$$\cong \operatorname{Hom}_{A_j}(\operatorname{Hom}_{A_j}(B_{1j}, A_j), A_j) \cong \operatorname{Hom}_{A_j}(B_{j1}, A_j),$$

which means that $S_1(\Lambda)$ is still a Frobenius-type triangular matrix algebra.

Using the same method in sequence, we can obtain the Frobenius-type triangular matrix algebra $S_k S_{k-1} \cdots S_1(\Lambda)$ for any k. In particular, it can be seen that $\Lambda \cong S_n S_{n-1} \cdots S_1(\Lambda)$.

A reflection functor F_1^+ : rep $(\Lambda) \to rep(S_1(\Lambda))$ can be described as follows. For

$$X = \begin{pmatrix} X_1 \\ X_2 \\ \vdots \\ X_n \end{pmatrix}_{\phi_{ij}} \in \operatorname{rep}(\Lambda),$$

define

$$F_1^+(X) = \begin{pmatrix} X_2 \\ \vdots \\ X_n \\ X_1' \end{pmatrix}_{\phi_{ij}'},$$

with $X'_1 = \operatorname{Ker}(X_{1, \operatorname{in}})$, where

$$X_{1, \text{ in}} : \bigoplus_{k=2}^{n} B_{1k} \bigotimes_{A_k} X_k \to X_1.$$

Denote by ψ_{i1} the composition of the inclusion map $X'_1 \hookrightarrow \bigoplus_{k=2}^n B_{1k} \bigotimes_{A_k} X_k$ and the projection $\bigoplus_{k=2}^n B_{1k} \bigotimes_{A_k} X_k \twoheadrightarrow B_{1i} \bigotimes_{A_i} X_i$. Then

$$\phi_{ij}' = \begin{cases} \overline{\psi_{i1}}, & \text{for } j = 1; \\ \phi_{ij}, & \text{otherwise.} \end{cases}$$

For a morphism $f = \{f_i\} : X \to Y$ in rep (Λ) , $F_1^+(f) = f' = \{f'_i\}$ where $f'_i = f_i$ for i = 2, ..., n and f'_1 is the unique morphism making the following diagram commutes:

Similarly, for

$$X = \begin{pmatrix} X_2 \\ \vdots \\ X_n \\ X_1 \end{pmatrix}_{\phi_{ij}} \in \operatorname{rep}(S_1(\Lambda)),$$

define $F_1^- : \operatorname{rep}(S_1(\Lambda)) \to \operatorname{rep}(\Lambda)$ satisfying

$$F_1^-(X) = \begin{pmatrix} X_1' \\ X_2 \\ \vdots \\ X_n \end{pmatrix}_{\phi_{ij}'}$$

with $X'_1 = \operatorname{Coker}(X_{1, \operatorname{out}})$, where $X_{1, \operatorname{out}} := (\overline{\phi_{j1}})_j : X_1 \to \bigoplus_{j=2}^n B_{1j} \bigotimes_{A_j} X_j$.

Denote by ψ_{1j} the composition of the inclusion map $B_{1j} \bigotimes_{A_j} X_j \hookrightarrow \bigoplus_{k=2}^n B_{1k} \bigotimes_{A_k} X_k$ and the projection $\bigoplus_{k=2}^n B_{1k} \bigotimes_{A_k} X_k \twoheadrightarrow X'_1$. Then

$$\phi_{ij}' = \begin{cases} \psi_{1j}, & \text{if } i = 1; \\ \phi_{ij}, & \text{otherwise.} \end{cases}$$

For a morphism $g = \{g_i\} : X \to Y$ in rep $(S_1(\Lambda))$, $F_1^-(g) = g' = \{g'_i\}$ where $g'_i = g_i$ for i = 2, ..., n and g'_1 is the unique morphism making the following diagram commutes:

$$\begin{array}{c} X_1 \xrightarrow{X_{1, \mathrm{out}}} \bigoplus_{k=2}^n B_{1k} \bigotimes_{A_k} X_k \longrightarrow \mathrm{Cok}(X_{1, \mathrm{out}}) \longrightarrow 0 \\ \downarrow^{g_1} \qquad \qquad \downarrow^{\oplus \mathrm{id} \otimes g_k} \qquad \qquad \stackrel{|g_1'|}{\stackrel{Y_1, \mathrm{out}}{\longrightarrow}} \bigoplus_{k=2}^n B_{1k} \bigotimes_{A_k} Y_k \longrightarrow \mathrm{Cok}(Y_{1, \mathrm{out}}) \longrightarrow 0. \end{array}$$

In the same way in sequence, for any k, we can define

$$F_k^+ : \operatorname{rep}(S_{k-1}S_{k-2}\dots S_1(\Lambda)) \to \operatorname{rep}(S_kS_{k-1}\dots S_1(\Lambda)),$$

$$F_k^- : \operatorname{rep}(S_kS_{k-1}\dots S_1(\Lambda)) \to \operatorname{rep}(S_{k-1}S_{k-2}\dots S_1(\Lambda)).$$

Then we denote

$$C^+ = F_n^+ F_{n-1}^+ \dots F_1^+ : \operatorname{rep}(\Lambda) \to \operatorname{rep}(\Lambda) \quad \text{and} \quad C^- = F_1^- F_2^- \dots F_n^- : \operatorname{rep}(\Lambda) \to \operatorname{rep}(\Lambda),$$

which are called the Coxeter functors on $\operatorname{rep}(\Lambda)$.

Remark 3.1 Let Q be a connected acyclic quiver. We can re-arrange the vertices of Q by making i < j if there is a path from j to i so as to get an admissible sequence 1, 2, ..., n with 1 as a sink vertex and n as a source vertex. Thus, the path algebra kQ can be seen as a special case of Frobenius-type triangular matrix algebras through assuming that $A_i = k$ for all i and B_{ij} is the k-linear spaces generated by all arrows from j to i. In this case, the reflection functors and Coxeter functors defined above accords with that as the classical case given by Dlab and Ringel in [6, 7, 18].

Proposition 3.2 For a Frobenius-type triangular matrix algebra Λ , and $X \in \operatorname{rep}(S_1(\Lambda)), Y \in \operatorname{rep}(\Lambda)$, there is a functorial isomorphism $\operatorname{Hom}_{\Lambda}(F_1^-(X), Y) \cong \operatorname{Hom}_{S_1(\Lambda)}(X, F_1^+(Y))$.

Proof Consider a morphism $f \in \text{Hom}_{S_1(\Lambda)}(X, F_1^+(Y))$. By definition, this is a collection of A_j -module homomorphisms $f_j : X_j \to (F_1^+(Y))_j$ with $1 \le j \le n$ satisfying certain commutative relations. In the diagram

$$\begin{array}{c} X_1 \xrightarrow{X_{1,\mathrm{out}}} \bigoplus_{k=2}^n B_{1k} \bigotimes_{A_k} X_k \longrightarrow \mathrm{Cok}(X_{1,\mathrm{out}}) \longrightarrow 0 \\ & \downarrow^{f_1} & \downarrow^{\oplus \mathrm{id} \otimes f_k} & \stackrel{|g_1}{\longrightarrow} \\ 0 \longrightarrow \mathrm{Ker}(Y_{1,\mathrm{in}}) \longrightarrow \bigoplus_{k=2}^n B_{1k} \bigotimes_{A_k} Y_k \xrightarrow{Y_{1,\mathrm{in}}} Y_1 \end{array}$$

its left square commutes. Observe that $Y_{1, \text{ in }} \circ (\oplus \text{id} \otimes f_k) \circ X_{1, \text{ out}} = 0$, so $Y_{1, \text{ in }} \circ (\oplus \text{id} \otimes f_k)$ factors through the cokernel of $X_{1, \text{ out}}$. Then, there is a map g_1 such that the right square commutes.

Thus if we set $g_j := f_j$ for $2 \le j \le n$, we get a homomorphism $g : F_1^-(X) \to Y$ corresponding to the given f. Write $\pi(f) = g$.

Conversely, consider a homomorphism $g: F_1^-(X) \to Y$, in the diagram

$$\begin{array}{c} X_{1} \xrightarrow{X_{1,\mathrm{out}}} \bigoplus_{k=2}^{n} B_{1k} \bigotimes_{A_{k}} X_{k} \longrightarrow \mathrm{Cok}(X_{1,\mathrm{out}}) \longrightarrow 0 \\ \downarrow & \downarrow \\ f_{1} & \downarrow \\ \forall & \downarrow \\ 0 \longrightarrow \mathrm{Ker}(Y_{1,\mathrm{in}}) \longrightarrow \bigoplus_{k=2}^{n} B_{1k} \bigotimes_{A_{k}} Y_{k} \xrightarrow{Y_{1,\mathrm{in}}} Y_{1} \end{array}$$

the right square commutes. There is a unique map f_1 such that the left square commutes.

Similarly as above, let $f_j := g_j$ for $2 \le j \le n$. Then we get a homomorphism $f : F_1^-(X) \to Y$ corresponding to the given g. Write $\tau(g) = f$.

For a morphism $f \in \text{Hom}_{S_1(\Lambda)}(X, F_1^+(Y))$, there exists a unique $\pi(f)$ makes the right square above commutes. And for $\pi(f)$, there exists a unique $\tau\pi(f)$ makes the left square above

commutes. Since f makes the left square above commutes, we have $\tau \pi(f) = f$. Similarly, $\pi \tau(g) = g$.

Therefore, π and τ are mutual-inverse, and we get a functorial isomorphism

$$\operatorname{Hom}_{\Lambda}(F_1^-(X), Y) \cong \operatorname{Hom}_{S_1(\Lambda)}(X, F_1^+(Y)).$$

Lemma 3.3 For a Frobenius-type triangular matrix algebra Λ , there is a short exact sequence of Λ - Λ -bimodules

$$P_{\bullet}: 0 \to \bigoplus_{1 \le i < j \le n} \Lambda e_i \bigotimes_{A_i} B_{ij} \bigotimes_{A_j} e_j \Lambda \xrightarrow{d} \bigoplus_{k=1}^n \Lambda e_k \bigotimes_{A_k} e_k \Lambda \xrightarrow{\text{mult}} \Lambda \to 0,$$

where d satisfies $d(p \otimes b \otimes q) := pb \otimes q - p \otimes bq$ and the morphism "mult" is given by the multiplication of Λ .

Proof Trivially, d is injective and mult is surjective. Also, Im(d) = Ker(mult).

Corollary 3.4 For a Frobenius-type triangular matrix algebra Λ and $X \in \operatorname{rep}_{l.f.}(\Lambda)$, there is a projective resolution of X:

$$P_{\bullet} \bigotimes_{\Lambda} X : 0 \to \bigoplus_{1 \le i < j \le n} P_i \bigotimes_{A_i} B_{ij} \bigotimes_{A_j} X_j \xrightarrow{d \otimes X} \bigoplus_{k=1}^n P_k \bigotimes_{A_k} X_k \xrightarrow{\text{mult}} X \to 0$$

with $(d \otimes X)(p \otimes b \otimes x) = pb \otimes x - p \otimes \phi_{ij}(b \otimes x).$

Proof Here P_i is just defined in (2.1). Then, $P_{\bullet} \bigotimes_{\Lambda} X$ is always exact. Since X is locally free, $e_k \Lambda \bigotimes_{\Lambda} X = e_k X$ are free A_k -modules. Thus $P_{\bullet} \bigotimes_{\Lambda} X$ is a projective resolution.

Following [9, 10], we define a functor T of rep(Λ) satisfying that

$$TX = \begin{pmatrix} X_1 \\ X_2 \\ \vdots \\ X_n \end{pmatrix}_{\psi_{ij}} \quad \text{for any } X = \begin{pmatrix} X_1 \\ X_2 \\ \vdots \\ X_n \end{pmatrix}_{\phi_{ij}} \in \operatorname{rep}(\Lambda),$$

where $\psi_{ij}(b \otimes x) = -\phi_{ij}(b \otimes x)$ for $b \in B_{ij}$, $x \in X_j$, $1 \le i < j \le n$. Obviously, T is an automorphism functor.

Theorem 3.5 For a Frobenius-type triangular matrix algebra Λ and each $X \in \operatorname{rep}_{l.f.}(\Lambda)$, there are functorial isomorphisms: $TC^+(X) \cong \tau(X)$ and $TC^-(X) \cong \tau^-(X)$.

This theorem and its preparation above follow the conclusion and method of [9, 10]. But, here Λ is a Frobenius-type triangular matrix algebra, which includes the classes of algebras in [9, 10]. We need to overcome the different key point such as the different expression form of algebras by using the dual basis in Lemma 2.4.

Firstly, from Λ , we construct a new Frobenius-type triangular matrix algebra:

$$\tilde{\Lambda} = \begin{pmatrix} \tilde{A}_1 & \tilde{A}_{1,2} & \dots & \tilde{A}_{1,2n} \\ 0 & \tilde{A}_2 & \dots & \tilde{A}_{2,2n} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \tilde{A}_{2n} \end{pmatrix},$$

where

$$\tilde{A}_i = \begin{cases} A_i, & \text{if } 1 \le i \le n, \\ A_{i-n}, & \text{if } n+1 \le i \le 2n, \end{cases}$$

and

$$\tilde{B}_{ij} = \begin{cases} B_{ij}, & \text{if } 1 \leq i < j \leq n \text{ or } n+1 \leq i < j \leq 2n, \\ \text{Hom}_{A_i}(B_{j-n,i}, A_i), & \text{if } 1 \leq j-n < i \leq n, \\ 0, & \text{otherwise}, \end{cases}$$

and $\tilde{A}_{ij} = \bigoplus_{l=0}^{j-i-1} \bigoplus_{i < k_1 < k_2 < \dots < k_l < j} \tilde{B}_{ik_1} \bigotimes_{\tilde{A}_{k_1}} \tilde{B}_{k_1k_2} \bigotimes \dots \bigotimes_{\tilde{A}_{k_l}} \tilde{B}_{k_lj}$ for $1 \le i < j \le 2n$. For any non-negative integer t, denote

$$1^{(t)} := \sum_{i=t+1}^{t+n} e_i, \quad 1_0^{(t)} := \sum_{i=1}^{t+n} e_i$$

and the corresponding subalgebras

$$\Lambda^{(t)} := 1^{(t)} \tilde{\Lambda} 1^{(t)}, \quad \tilde{\Lambda}^{(t)} := 1_0^{(t)} \tilde{\Lambda} 1_0^{(t)}$$

for $0 \le t \le n$. It is easy to see that $\tilde{\Lambda}^{(0)} \cong \Lambda^{(0)} \cong \Lambda^{(n)} \cong \Lambda$, $\tilde{\Lambda}^{(n)} \cong \tilde{\Lambda}$. And $\Lambda^{(t)} \cong S_t \cdots S_1(\Lambda)$ for $1 \le t \le n$.

Let

$$X = \begin{pmatrix} X_1 \\ \vdots \\ X_{2n} \end{pmatrix}_{\phi_{ij}} \in \operatorname{rep}(\tilde{\Lambda})$$

satisfy the following condition:

$$X_{n+i} \xrightarrow{\tilde{X}_{i,\text{ out}}} \bigoplus_{k=i+1}^{n+i-1} B_{ik} \bigotimes_{A_k} X_k \xrightarrow{\tilde{X}_{i,\text{ in}}} X_i \text{ such that } \tilde{X}_{i,\text{ in}} \circ \tilde{X}_{i,\text{ out}} = 0 \quad \text{for } 1 \le i \le n.$$
(3.1)

We define the restriction functors:

$$\operatorname{Res}^{(t,m)} : \operatorname{rep}(\tilde{\Lambda}^{(m)}) \to \operatorname{rep}(\Lambda^{(t)}), \quad X \mapsto 1^{(t)} \tilde{\Lambda}^{(m)} \otimes_{\tilde{\Lambda}^{(m)}} X;$$

$$\operatorname{Res}_{(t,m)} : \operatorname{rep}(\tilde{\Lambda}^{(m)}) \to \operatorname{rep}(\tilde{\Lambda}^{(t)}), \quad X \mapsto 1^{(t)}_{0} \tilde{\Lambda}^{(m)} \otimes_{\tilde{\Lambda}^{(m)}} X \quad \text{for } 1 \le t \le m \le n.$$

Obviously, $\operatorname{Res}_{(t,m)}$ has a right adjoint $\operatorname{Res}_{(t,m)}^*(-) = \operatorname{Hom}_{\tilde{\Lambda}^{(t)}}(1_0^{(t)}\tilde{\Lambda}^{(m)}, -).$

The following lemma is a generalization of [10, Lemma 10.2]. Their proofs are identical.

Lemma 3.6 With the above notations, there is a functorial isomorphisms

$$\operatorname{Res}^{(i,i)} \circ \operatorname{Res}^{*}_{(i-1,i)}(X) \cong F_{i}^{+} \circ \operatorname{Res}^{(i-1,i-1)}(X)$$

for $X \in \operatorname{rep}(\tilde{\Lambda}^{(i-1)})$ which satisfies (3.1) for all $1 \leq i \leq n$.

By Lemma 3.6, we obtain that for any $X \in \operatorname{rep}(\Lambda)$ (regarded as a representation of $\operatorname{rep}(\tilde{\Lambda}^{(0)})$), we have

$$\operatorname{Hom}_{\Lambda}(1^{(0)}\Lambda 1^{(n)}, X) = \operatorname{Res}^{(n,n)} \circ \operatorname{Res}^{*}_{(n-1,n)} \circ \cdots \circ \operatorname{Res}^{*}_{(0,1)}(X)$$
$$= F_{n}^{+} \circ \operatorname{Res}^{n-1} \circ \operatorname{Res}^{*}_{(n-2,n-1)} \circ \cdots \circ \operatorname{Res}^{*}_{(0,1)}(X)$$
$$= \cdots$$

Representations of Frobenius-type Triangular Matrix Algebras

$$= F_n^+ \circ F_{n-1}^+ \circ \cdots F_1^+ \circ \operatorname{Res}^0(X)$$
$$= C^+(X).$$

Denote

 $\operatorname{Res}_{0} = \operatorname{Res}_{(0,1)} \circ \operatorname{Res}_{(1,2)} \circ \cdots \circ \operatorname{Res}_{(n-1,n)}, \quad \operatorname{Res}_{0}^{*} = \operatorname{Res}_{(n-1,n)}^{*} \circ \cdots \circ \operatorname{Res}_{(1,2)}^{*} \circ \operatorname{Res}_{(0,1)}^{*}.$

It is easy to see that Res_0^* is right adjoint of Res_0 .

Now, following [9, 10], we construct another functor $R_0^* : \operatorname{rep}(\Lambda^{(0)}) \to \operatorname{rep}(\tilde{\Lambda})$, and then show that R_0^* is right adjoint to Res_0 .

Let $X \in \operatorname{rep}(\Lambda^{(0)})$. We first define $\tilde{X} \in \operatorname{rep}(\tilde{\Lambda})$ by requiring that

$$\operatorname{Res}^{(0,n)}(\tilde{X}) = X,$$

$$\operatorname{Res}^{(n,n)}(\tilde{X}) = \bigoplus_{1 \le k < t \le n} \operatorname{Hom}_{A_t}\left(\tilde{B}_{t,n+k} \bigotimes_{A_k} e_{n+k}\Lambda^{(n)}, X_t\right).$$

For $1 \le i < j \le n$, it remains to define the structure map of \tilde{X} as A_j -module morphisms:

$$\phi_{j,n+i}: \tilde{B}_{j,n+i} \bigotimes_{A_i} \tilde{X}_{n+i} \to \tilde{X}_j = X_j,$$

which is given by the following composition:

1

$$\begin{split} \tilde{B}_{j,n+i} \bigotimes_{A_i} \left(\bigoplus_{1 \le k < t \le n} \operatorname{Hom}_{A_t} \left(\tilde{B}_{t,n+k} \bigotimes_{A_K} e_{n+k} \Lambda^{(n)} e_{n+i}, X_t \right) \right) \\ \xrightarrow{\operatorname{proj.}} \tilde{B}_{j,n+i} \bigotimes_{A_i} \operatorname{Hom}_{A_j} \left(\tilde{B}_{j,n+i} \bigotimes_{A_i} e_{n+i} \Lambda^{(n)} e_{n+i}, X_j \right) \\ &= \tilde{B}_{j,n+i} \bigotimes_{A_i} \operatorname{Hom}_{A_j} \left(\tilde{B}_{j,n+i}, X_j \right) \\ \xrightarrow{\operatorname{eval.}} X_j, \end{split}$$

where the first map is the projection on the direct summand indexed by i, j and the second map is the evaluation $b \otimes \varphi = \varphi(b)$.

Secondly, we define a $\tilde{\Lambda}$ -subrepresentation $R_0^*(X)$ of \tilde{X} as follows. We set

$$(R_0^*(X))_i = \tilde{X}_i = X_i, \quad 1 \le i \le n,$$

and define $R_0^*(X)_{n+i}$ as the subspace of \tilde{X}_{n+i} generated by all of

$$(\mu_{k,t}^i)_{1 \le k < t \le n} \in \bigoplus_{1 \le k < t \le n} \operatorname{Hom}_{A_t} \left(\tilde{B}_{t,n+k} \bigotimes_{A_k} e_{n+k} \Lambda^{(n)} e_{n+i}, X_t \right)$$

such that, for all $1 \le t \le n$ and $\lambda \in e_{n+t} \Lambda^{(n)} e_{n+i}$, the following relation holds:

$$\sum_{\leq k < t, \ell \in L_{kt}} \mu^i_{k,t}(\ell^* \otimes \ell \lambda) + \sum_{t < m \leq n, r \in R_{tm}} \phi_{tm}(r \otimes \mu^i_{t,m}(r^* \otimes \lambda)) = 0.$$
(3.2)

For $\mu^j = (\mu_{k,t}^j)_{1 \le k < t \le n} \in (R_0^*(X))_{n+j}, 1 \le j \le n$, we deduce from the definitions that $\tilde{X} := \circ \tilde{X} := (\mu^j)$

$$= \tilde{X}_{j,\mathrm{in}} \left(\sum_{1 \le i < j, \ell \in L_{ij}} (\ell^* \otimes \phi_{n+i,n+j}(\ell \otimes \mu^j)) + \sum_{j < k \le n, r \in R_{ij}} (r \otimes \phi_{k,n+j}(r^* \otimes \mu^j)) \right)$$

Li F. and Ye C.

$$= \sum_{1 \le i < j, \ell \in L_{ij}} \phi_{j,n+i}(\ell^* \otimes \phi_{n+i,n+j}(\ell \otimes \mu^j)) + \sum_{j < k \le n, r \in R_{ij}} \phi_{j,k}(r \otimes \phi_{k,n+j}(r^* \otimes \mu^j))$$
$$= \sum_{1 \le i < j, \ell \in L_{ij}} \mu_{i,j}^j(\ell^* \otimes \ell \otimes e_{n+j}) + \sum_{j < k \le n, r \in R_{ij}} \phi_{j,k}(r \otimes \mu_{j,k}^j(r^* \otimes e_{n+j})) = 0.$$

So, $R_0^*(X)$ satisfies (3.1).

Thus, we have obtained a functor $R_0^* : \operatorname{rep}(\Lambda^{(0)}) \to \operatorname{rep}(\tilde{\Lambda}); X \mapsto R_0^*(X)$. We have that R_0^* is isomorphic to Res_0^* . This is according to the uniqueness of adjoint functors up to natural equivalence and the following lemma.

Lemma 3.7 R_0^* is right adjoint to Res_0 .

It is a generalization of [10, Lemma 10.3]. Their proofs are identical.

Lemma 3.8 Let A_i be a Frobenius algebra, U and V be finitely generated free A_i -modules. Then we have an isomorphism $D\operatorname{Hom}_{A_i}(U,V) \cong \operatorname{Hom}_{A_i}(V,U)$.

Proof Since A_i is a Frobenius algebra, $DA_i \cong A_i$. Let $U = \bigoplus_{j=1}^n A_i$ and $V = \bigoplus_{t=1}^m A_i$. Then $D\operatorname{Hom}_{A_i}(U,V) \cong \bigoplus_{j=1}^n \bigoplus_{t=1}^m D\operatorname{Hom}_{A_i}(A_i,A_i) \cong \bigoplus_{j=1}^n \bigoplus_{t=1}^m \operatorname{Hom}_{A_i}(A_i,A_i) \cong \operatorname{Hom}_{A_i}(V,U)$. **Proposition 3.9** For a Frobenius-type triangular matrix algebra Λ and $X \in \operatorname{rep}_{l.f.}(\Lambda)$, there is an isomorphism $\tau(TM) \cong \operatorname{Res}^{(n,n)} \circ R_0^*(X)$, where Λ is identified with $\Lambda^{(0)}$ and $\Lambda^{(n)}$ by their

definitions.

Proof The proof is similar to [10, Proposition 10.4]. The only illustration we need to add is the fact that for a locally free Λ -module X, we have $D\operatorname{Hom}_{A_i}(e_i\Lambda, X_i) \cong \operatorname{Hom}_{A_i}(X_i, e_i\Lambda)$. This follows from Lemma 3.8.

Proof of Theorem 3.5 By Proposition 3.9, if $X \in \operatorname{rep}_{l.f.}(\Lambda)$, we have

$$TC^+(X) \cong \operatorname{Res}^{(n,n)} \circ \operatorname{Res}^*_0(TX) \cong \operatorname{Res}^{(n,n)} \circ R^*_0(TX) \cong \tau(T^2X) \cong \tau(X).$$

Let $X, Y \in \operatorname{rep}_{l, f}(\Lambda)$. Then we have

$$\operatorname{Hom}_{\Lambda}(\tau^{-}(X), Y) \cong \operatorname{Hom}_{\Lambda}(X, \tau(Y))$$
$$\cong \operatorname{Hom}_{\Lambda}(X, C^{+}(TY))$$
$$\cong \operatorname{Hom}_{\Lambda}(C^{-}(TX), Y)$$

The first isomorphism is obtained by Corollary 2.8 and the third isomorphism follows from Proposition 3.2.

There exists a functorial isomorphism of right Λ -modules $DX \cong \operatorname{Hom}_{\Lambda}(X, D\Lambda)$ for all Λ -modules X. Since $D\Lambda \in \operatorname{rep}_{l.f.}(\Lambda)$, taking $Y = D\Lambda$ we get $\tau^{-}(X) \cong C^{-}(TX)$.

Recall that the category of Gorenstein-projective modules of Λ is

$$\mathcal{GP}(\Lambda) = \{ X \in \operatorname{rep}(\Lambda) | \operatorname{Ext}^{1}_{\Lambda}(X, \Lambda) \} = 0.$$

As an immediate consequence of Theorem 3.5 and the definition of $C^+(-)$, we get the following result.

Corollary 3.10 For a Frobenius-type triangular matrix algebra Λ and $X \in \operatorname{rep}_{l.f.}(\Lambda)$, the following are equivalent:

(i) $X \in \mathcal{GP}(\Lambda);$ (ii) $C^+(X) = 0;$ (iii) $X_{i, \text{ in }}$ is injective for all $1 \leq i \leq n$.

This result has been proved in a more general case in [15].

4 Root Systems in Case of Dynkin Type

For a Frobenius-type triangular matrix algebra Λ , let $C = (c_{ij}) \in M_n(\mathbb{Z})$, where

$$c_{ij} = \begin{cases} 2, & \text{if } i = j, \\ -\text{rank}_{A_i}(B_{ij}), & \text{if } i < j, \\ -\text{rank}_{A_i}(B_{ji}), & \text{if } i > j. \end{cases}$$

Denote $c_i = \dim_k(A_i)$. Then it is easy to get that $c_i c_{ij} = c_j c_{ji} = -\dim_k(B_{ij})$, which means that C is a symmetrizable Cartan matrix.

Define a quadratic form $q_C : \mathbb{Z}^n \to \mathbb{Z}$ of C satisfying for $x = (x_1, \ldots, x_n)^t \in \mathbb{Z}^n$,

$$q_C(x) := \sum_{i=1}^n c_i x_i^2 - \sum_{i < j} c_i |c_{ij}| x_i x_j.$$
(4.1)

The Cartan matrix C is said to be of Dynkin type (resp. Euclidean type) if q_C is positive define (resp. positive semidefinite).

We define the valued quiver $\Gamma(\Lambda)$ via the Cartan matrix C whose vertices are $1, \ldots, n$ and whose arrow $i \leftarrow j$ from j to i exists for each pair (i, j) with i < j if $c_{ij} < 0$, with valuation $(-c_{ji}, -c_{ij})$ on the arrow $i \leftarrow j$. It is well known, see [12, Theorem 4.8], that

Fact C is of Dynkin type if and only if $\Gamma(\Lambda)$ is a disjoint union of quivers whose underlying valued graph is of Dynkin type.

The standard basis vectors of \mathbb{Z}^n , denoted as $\alpha_1, \ldots, \alpha_n$, are the positive simple roots of the Kac–Moody algebra $\mathfrak{g}(C)$ associated with C, that is, of the quadratic form $q_C(x)$. For $1 \leq i, j \leq n$, define the reflections s_i satisfying that $s_i(\alpha_j) := \alpha_j - c_{ij}\alpha_i$.

The Weyl group W(C) of $\mathfrak{g}(C)$ is the subgroup of $\operatorname{Aut}(\mathbb{Z}^n)$ generated by s_1, \ldots, s_n . It is well known that W(C) is finite if and only if C is of Dynkin type.

Let $\Delta(C)$ be the set of roots of C and $\Delta_{re}(C) := \bigcup_{i=1}^{n} W(\alpha_i)$ be the set of *real roots* of C. Let $\Delta^+(C) := \Delta(C) \cap \mathbb{N}^n$ and $\Delta_{re}^+(C) := \Delta_{re}(C) \cap \mathbb{N}^n$. The following are equivalent:

(i) C is of Dynkin type;

(ii) $\Delta(C)$ is finite;

(iii) $\Delta_{\rm re}(C) = \Delta(C)$. Let

$$\beta_k = \begin{cases} \alpha_1, & \text{if } k = 1, \\ s_1 s_2 \cdots s_{k-1}(\alpha_k), & \text{if } 2 \le k \le n \end{cases} \text{ and } \gamma_k = \begin{cases} s_n s_{n-1} \cdots s_{k+1}(\alpha_k), & \text{if } 1 \le k \le n-1, \\ \alpha_n, & \text{if } k = n. \end{cases}$$

Let $c^+ = s_n s_{n-1} \cdots s_1 : \mathbb{Z}^n \to \mathbb{Z}^n$ and $c^- = s_1 s_2 \cdots s_n : \mathbb{Z}^n \to \mathbb{Z}^n$ be the Coxeter transformations. For $k \in \mathbb{Z}$, set

$$c^{k} := \begin{cases} (c^{+})^{k}, & \text{if } k > 0, \\ (c^{-})^{-k}, & \text{if } k < 0, \\ \text{id}, & \text{if } k = 0. \end{cases}$$

It follows that $c^+(\beta_k) = -\gamma_k$.

The following three lemmas are well known, for example see [3, Chapter VII].

Lemma 4.1 Suppose that C is not of Dynkin type. Then the element $c^{-r}(\beta_i)$ and $c^s(\gamma_j)$ with $r, s \ge 0$ and $1 \le i, j \le n$ are pairwise different elements in $\Delta_{re}^+(C)$.

Let C be of Dynkin type. Let $p_i \ge 1$ be minimal with $c^{-p_i}(\beta_i) \notin \mathbb{N}^n$ for $1 \le i \le n$, and let $q_j \ge 1$ be minimal with $c^{q_j}(\gamma_j) \notin \mathbb{N}^n$ for $1 \le j \le n$. The elements $c^{-r}(\beta_i)$ with $1 \le i \le n$ and $0 \le r \le p_i - 1$ are pairwise different, and the elements $c^s(\gamma_j)$ with $1 \le j \le n$ and $0 \le s \le q_j - 1$ are pairwise different.

Lemma 4.2 Assume that C is of Dynkin type. Then $\Delta^+(C) = \{c^{-r}(\beta_i) | 1 \le i \le n, 0 \le r \le p_i - 1\} = \{c^s(\gamma_j) | 1 \le j \le n, 0 \le s \le q_j - 1\}.$

Lemma 4.3 Assume that C is of Dynkin type. For every positive vector \underline{x} , there exist $s \ge 0$ such that $c^{s}\underline{x} > 0$ but $c^{s+1}\underline{x} \ge 0$, and $t \ge 0$ such that $c^{-t}\underline{x} > 0$ but $c^{-t-1}\underline{x} \ge 0$.

Definition 4.4 For a Frobenius-type triangular matrix algebra Λ , let X be a locally free Λ module. Let r_i be the rank of the free A_i -module X_i . Following [10], we denote

$$\underline{\operatorname{rank}}(X) := (r_1, \dots, r_n).$$

Lemma 4.5 $\underline{\operatorname{rank}}(P_k) = \beta_k, \ \underline{\operatorname{rank}}(I_k) = \gamma_k.$

This lemma is a generalization of [10, Lemmas 3.2 and 3.3]. Its proof is similar to that of [10, Lemma 3.2], using the resolution (2.2) and (2.4).

Definition 4.6 For a Frobenius-type triangular matrix algebra Λ , following [10], a Λ -module X is called τ -locally free if $\tau^k(X)$ is locally free for all $k \in \mathbb{Z}$. Moreover, X is called indecomposable τ -locally free if X cannot be written as a sum of two proper τ -locally free Λ -modules.

The following proposition is a generalization of [10, Propositions 9.6 and 11.4].

Proposition 4.7 For a Frobenius-type triangular matrix algebra Λ , let X be a rigid and locally free Λ -module. Then

(i) $F_1^+(X)$ is a rigid and locally free $S_1(\Lambda)$ -module and $F_n^-(X)$ is a rigid and locally free $S_{n-1} \cdots S_1(\Lambda)$ -module;

(ii) X is τ -locally free and $\tau^k(X)$ is rigid for all $k \in \mathbb{Z}$.

Proof Its proof is identical to the proof of [10, Propositions 9.6 and 11.4]. The only illustration we need to add is the fact that for a locally free Λ -module X and its minimal projective resolution of the form $0 \to P'' \to P' \to X \to 0$, we have that P' and P'' are two direct sums of P_i for $1 \leq i \leq n$. The conclusion follows from Proposition 2.6, Corollary 2.8 and the horseshoe lemma.

Lemma 4.8 For a Frobenius-type triangular matrix algebra Λ , let X be an indecomposable τ -locally free Λ -module. Then X is isomorphic to E_1 if and only if $F_1^+(X) = 0$ (or equivalently, $s_1(\underline{\operatorname{rank}}(X)) \neq 0$). If $X \ncong E_1$, then $F_1^+(X)$ is an indecomposable τ -locally free $S_1(\Lambda)$ -module and $\underline{\operatorname{rank}}(F_1^+(X)) = s_1(\underline{\operatorname{rank}}(X))$.

Proof Since X is indecomposable τ -locally free, we obtain that $X_{1, \text{ in}}$ is surjective. If $X \ncong E_1$, we have an exact sequence $0 \longrightarrow \text{Ker}(X_{1, \text{ in}}) \longrightarrow \bigoplus_{k=2}^n B_{1k} \bigotimes_{A_k} X_k \longrightarrow X_1 \longrightarrow 0$. So

$$(\underline{\operatorname{rank}}(F_1^+(X)))_1 = \sum_{k=2}^n |c_{1k}| a_k - a_1 = (s_1(\underline{\operatorname{rank}}(X)))_1$$

If $X \cong E_1$, we have $F_1^+(X) = 0$ by the definition of F_1^+ , and $s_1(\underline{\operatorname{rank}}(E_1))_1 = -1$.

We need the following two propositions, which are generalizations of [10, Propositions 11.5 and 11.6] respectively. Their proofs are similar.

Proposition 4.9 For a Frobenius-type triangular matrix algebra Λ and an indecomposable τ -locally free Λ -module X, the following statements hold:

(i) If $\tau^k(X) \neq 0$ for some $k \in \mathbb{Z}$, then $\underline{\operatorname{rank}}(\tau^k(X)) = c^k(\underline{\operatorname{rank}}(X));$

(ii) If $\tau^k(X) \neq 0$ for some $k \in \mathbb{Z}$ and $\underline{\operatorname{rank}}(X)$ is contained in $\Delta^+(C)$, then $\underline{\operatorname{rank}}(\tau^k(X))$ is in $\Delta^+(C)$.

Proposition 4.10 For a Frobenius-type triangular matrix algebra Λ and a Λ -module X, if either $X \cong \tau^{-k}(P_i)$ or $X \cong \tau^k(I_i)$ for some $k \ge 0$ and $1 \le i \le n$, the following statements hold:

(i) X is τ -locally free and rigid;

(ii) $\underline{\operatorname{rank}}(X) \in \Delta_{\operatorname{re}}^+(C);$

(iii) If either $Y \cong \tau^{-m}(P_j)$ or $Y \cong \tau^m(I_j)$ for some $m \ge 0$ and $1 \le j \le n$ with $\underline{\operatorname{rank}}(X) = \underline{\operatorname{rank}}(Y)$, then $X \cong Y$.

Theorem 4.11 For a Frobenius-type triangular matrix algebra Λ , the following statements hold:

(a) The number of isomorphism classes of indecomposable τ -locally free Λ -modules is finite if and only if C is of Dynkin type.

(b) If C is of Dynkin type, then the mapping $\underline{\operatorname{rank}} : X \mapsto \underline{\operatorname{rank}}(X)$ induces a bijection between the set of isomorphism classes of indecomposable τ -locally free Λ -modules and the set of positive roots of the quadratic form $q_C(x)$.

Proof In the case that C is not of Dynkin type, by Lemmas 4.1, 4.5 and Proposition 4.9, we know there are infinitely many isomorphism classes of indecomposable τ -locally free Λ -modules.

In the case that C is of Dynkin type, firstly, we need to prove that $\underline{\operatorname{rank}}(X) \in \Delta^+(C)$ for any indecomposable τ -locally free Λ -module X. We denote $\underline{\operatorname{rank}}(X) = \underline{x}$. By Lemma 4.3, there exists a least s such that $c^s \underline{x} > 0$ but $c^{s+1} \underline{x} \neq 0$. Because $c^+ = s_n \cdots s_1$, there also exists a least i such that $0 \leq i \leq n-1$, $s_i \cdots s_1 c^t \underline{x} > 0$, but $s_{i+1} \cdots s_1 c^t \underline{x} \neq 0$.

We know that $X' = F_i^+ \cdots F_1^+ C^{+t} X$ is indecomposable τ -locally free by Lemma 4.8 and that

$$\underline{\operatorname{rank}}(F_i^+ \cdots F_1^+ C^{+t} X) = s_i \cdots s_1 c^t \underline{x}.$$

Because $s_{i+1}(\underline{\operatorname{rank}}(X')) \neq 0$, there is an isomorphism $X' \cong E_{i+1}$ by Lemma 4.8. So $s_i \cdots s_1 c^t \underline{x} = \alpha_{i+1}$, and according to Lemma 4.2, the vector $\underline{x} = c^{-t} s_1 \cdots s_i \alpha_{i+1} = c^{-t} \beta_{i+1}$ is a positive root of $q_C(x)$ and $\underline{\operatorname{rank}}(-)$ is surjective.

If X and Y are indecomposable τ -locally free Λ -modules such that $\underline{\operatorname{rank}}(X) = \underline{\operatorname{rank}}(Y)$, then we have, as earlier

$$F_i^+ \cdots F_1^+ C^{+t} X \cong E_{i+1} \cong F_i^+ \cdots F_1^+ C^{+t} Y,$$

so that

$$X \cong C^{-t}F_1^- \cdots F_i^- E_{i+1} \cong Y$$

Thus $\underline{\operatorname{rank}}(-)$ is an injective mapping.

Corollary 4.12 For the path algebra $\Lambda = AQ$ of an acyclic quiver Q over a Frobenius algebra A, the following statements hold:

(a) The number of isomorphism classes of indecomposable τ -locally free Λ -modules is finite if and only if Q is of Dynkin type;

(b) In the case that Q is of Dynkin type, the mapping $\underline{\operatorname{rank}} : X \mapsto \underline{\operatorname{rank}}(X)$ induces a bijection between the set of isomorphism classes of indecomposable τ -locally free Λ -modules and the set of positive roots of the quadratic form $q_Q(x) = \sum_{i \in Q_0} x_i^2 - \sum_{\alpha \in Q_1} x_{s(\alpha)} x_{t(\alpha)}$, where $x = (x_1, \ldots, x_n)^t \in \mathbb{Z}^n$.

Proof By Remark 1.2 (iii), $\Lambda = AQ$ is a Frobenius-type triangular matrix algebra via taking all $A_i = A$ and $B_{ij} = \bigoplus_{s=1}^{\#\{\alpha: j \to i\}} A$. So, $c_i = c_j = \dim_k A$, and then

$$-c_{ij} = -c_{ji} = \dim_k B_{ij} / \dim_k A = \#\{\alpha : j \to i\},\$$

which means that the Cartan matrix C is symmetric and in its corresponding valued quiver $\Gamma(\Lambda)$, the valuation $(-c_{ij}, -c_{ji})$ is given with the number of arrows from j to i in Q. By the definitions of the quadratic forms, we have

$$q_C(x) = \sum_{i=1}^n c_i x_i^2 - \sum_{i < j} c_i |c_{ij}| x_i x_j = \dim_k A\left(\sum_{i=1}^n x_i^2 - \sum_{i < j} |c_{ij}| x_i x_j\right) = q_Q(x) \dim_k A.$$

Hence, the positivity definite property of q_C and q_Q are the same with each other. Thus, the statements (a) and (b) follow respectively from Theorem 4.11 (a) and (b).

Recall in [14] for a generalized path algebra $\Lambda = k(Q, \mathcal{A})$, there is a corresponding valued quiver $\Upsilon(\Lambda)$. The set of vertices $\Upsilon(\Lambda)_0 = Q_0$. If there are arrows from j to i in Q, we give an arrow $i \leftarrow j$ in $\Upsilon(\Lambda)$, with valuation (d_{ji}, d_{ij}) where $d_{ji} = |Q_{ij}| \dim_k A_i$ and $d_{ij} = |Q_{ij}| \dim_k A_j$. Here $|Q_{ij}|$ means the number of arrows.

In the case when Q is acyclic, in order to realize $k(Q, \mathcal{A})$ as a triangular matrix algebra, we re-arrange the order of vertices in Q via assuming i < j if there exists a path from j to i. Let $B_{ij} = A_i Q_{ij} A_j$ for Q_{ij} the set of arrows from j to i in Q and then define A_{ij} as in Definition 1.1 and put A_i at the (i, i)-array. Then we obtain the triangular matrix algebra which is equal to $\Lambda = k(Q, \mathcal{A}).$

Moreover, $-c_{ij} = \operatorname{rank}_{A_i}(B_{ij}) = |Q_{ij}| \dim_k A_j = d_{ij}$. Thus, it follows that the valued quiver $\Upsilon(\Lambda)$ is coincident with $\Gamma(\Lambda)$.

Define a quadratic form $q_{k(Q,\mathcal{A})}: \mathbb{Z}^n \to \mathbb{Z}$ of $k(Q,\mathcal{A})$ satisfying for $x = (x_1, \ldots, x_n)^t \in \mathbb{Z}^n$,

$$q_{k(Q,\mathcal{A})}(x) = \sum_{i \in Q_0} d_i x_i^2 - \sum_{\alpha \in Q_1} d_{s(\alpha)} d_{s(\alpha)t(\alpha)} x_{s(\alpha)} x_{t(\alpha)}.$$
(4.2)

Comparing this quadratic form with that in (4.1), it is easy to see that $q_{k(Q,\mathcal{A})}$ is the special case of q_C for $\Lambda = k(Q, \mathcal{A})$.

Corollary 4.13 For an acyclic quiver Q and its generalized path algebra $\Lambda = k(Q, \mathcal{A})$ endowed by Frobenius algebras A_i at all vertices $i \in Q_0$, the following statements hold:

(a) The number of isomorphism classes of indecomposable τ -locally free Λ -modules is finite if and only if $\Upsilon(\Lambda)$ is of Dynkin type.

(b) If $\Omega(\Lambda)$ is of Dynkin type, then the mapping $\underline{\operatorname{rank}} : X \mapsto \underline{\operatorname{rank}}(X)$ induces a bijection between the set of isomorphism classes of indecomposable τ -locally free Λ -modules and

the set of positive roots of the $q_{k(Q,\mathcal{A})}(x) = \sum_{i \in Q_0} d_i x_i^2 - \sum_{\alpha \in Q_1} d_{s(\alpha)} d_{t(\alpha)} x_{s(\alpha)} x_{t(\alpha)}$, where $x = (x_1, \ldots, x_n)^t \in \mathbb{Z}^n$ and $d_i = \dim(A_i)$.

Proof They follow directly from $\Upsilon(\Lambda) = \Gamma(\Lambda)$, the Fact, Theorem 4.11 and that $q_{k(Q,\mathcal{A})} = q_C$ for $\Lambda = k(Q,\mathcal{A})$.

5 Analog of APR-tilting Module for Λ

The APR-tilting modules were introduced by Auslander et al. in [4] to interpret BGP-reflection functors as homomorphism functors of certain tilting modules. Also, it was the beginning of tilting theory.

For a Frobenius-type triangular matrix algebra Λ , for the case i = 1, we define $T_1 := \Lambda/P_1 \bigoplus \tau^-(P_1)$ and call T_1 a generalized APR-tilting module of Λ . This case follows from the fact that i = 1 is a "sink vertex" so as to gain the reflection functor.

Proposition 5.1 For a Frobenius-type triangular matrix algebra Λ , T_1 is a tilting Λ -module.

Proof Since P_1 is a rigid and locally free, $\tau^-(P_1)$ is rigid and locally free by Proposition 4.7. So T_1 is locally free. Then by Corollary 2.8, proj.dim $T_1 \leq 1$. Thus, $\operatorname{Ext}^1_{\Lambda}(T_1, T_1) \cong D\operatorname{Hom}_{\Lambda}(T_1, \tau(T_1)) = D\operatorname{Hom}_{\Lambda}(T_1, P_1) = 0$. Because Λ is connected, P_1 has no injective summand. Since τ^- takes non-injective indecomposable modules to non-projective indecomposable modules, $T_1 := \Lambda/P_1 \bigoplus \tau^-(P_1)$ has the same number of summands as primitive idempotents of Λ . So T_1 is a tilting Λ -module.

Remark 5.2 A similar analog of generalized APR-tilting modules was introduced in [16] for a class of triangular matrix algebras.

Lemma 5.3 For a Frobenius-type triangular matrix algebra Λ , there is an algebra isomorphism $End_{\Lambda}(T_1) \cong S_1(\Lambda).$

Proof Clearly, when $2 \le i, j \le n$,

$$e'_{i} \operatorname{End}_{\Lambda}(T_{1})e'_{j} \cong \operatorname{Hom}_{\operatorname{End}(T_{1})}(\operatorname{End}_{\Lambda}(T_{1})e'_{i}, \operatorname{End}_{\Lambda}(T_{1})e'_{j})$$
$$\cong \operatorname{Hom}_{\operatorname{End}(T_{1})}(\operatorname{Hom}(T_{1}, P_{i}), \operatorname{Hom}(T_{1}, P_{j}))$$
$$\cong \operatorname{Hom}_{\Lambda}(P_{i}, P_{j})$$
$$\cong e_{i}\Lambda e_{j}.$$

When i = 1, j = 1, $e'_i \operatorname{End}_{\Lambda}(T_1)e'_j \cong \operatorname{Hom}_{\Lambda}(\tau^-(P_1), \tau^-(P_1)) \cong \operatorname{Hom}_{\Lambda}(P_1, P_1) \cong A_1$. When i = 1, j > 1, $e'_i \operatorname{End}_{\Lambda}(T_1)e'_j \cong \operatorname{Hom}_{\Lambda}(\tau^-(P_1), P_j) = 0$. When j = 1, i > 1, $e'_i \operatorname{End}_{\Lambda}(T_1)e'_j \cong \operatorname{Hom}_{\Lambda}(P_i, \tau^-(P_1))$.

Since there is a minimal injective resolution:

$$0 \to P_1 \to I_1 \to \bigoplus_{j=2}^n I_j \bigotimes_{A_j} B_{j1} \to 0,$$

then we have

$$0 \to \nu^{-}(P_1) \to P_1 \to \bigoplus_{j=2}^n P_j \otimes_{A_j} B_{j1} \to \tau^{-}(P_1) \to 0.$$

Apply functor $\operatorname{Hom}_{\Lambda}(P_i, -)$ for i > 1.

Since $\operatorname{Hom}_{\Lambda}(P_i, P_1) = 0$, $\operatorname{Hom}_{\Lambda}(P_i, \bigoplus_{k=2}^n P_k \bigotimes_{A_k} B_{k1}) \to \operatorname{Hom}_{\Lambda}(P_i, \tau^-(P_1))$ is injective.

Since P_i is projective Λ -module, $\operatorname{Hom}_{\Lambda}(P_i, \bigoplus_{k=2}^n P_k \bigotimes_{A_k} B_{k1}) \to \operatorname{Hom}_{\Lambda}(P_i, \tau^-(P_1))$ is surjective.

So, $\operatorname{Hom}_{\Lambda}(P_i, \bigoplus_{k=2}^n P_k \bigotimes_{A_k} B_{k1}) \cong \operatorname{Hom}_{\Lambda}(P_i, \tau^-(P_1)).$ Then, $e'_i \operatorname{End}_{\Lambda}(T_1) e'_1 \cong \operatorname{Hom}_{\Lambda}(P_i, \bigoplus_{k=2}^n P_k \bigotimes_{A_k} B_{k1}) \cong \bigoplus_{k=2}^n e_1 \Lambda e_k \bigotimes_{A_k} B_{k1} \cong A_{i1}.$ At last, $\operatorname{End}_{\Lambda}(T_1) \cong S_1(\Lambda).$

Using Proposition 5.1 and Lemma 5.3, we can prove the following theorem.

Theorem 5.4 For a Frobenius-type triangular matrix algebra Λ , there is a functorial isomorphism

$$F_1^+(-) \cong \operatorname{Hom}_{\Lambda}(T_1,-) : \operatorname{rep}(\Lambda) \to \operatorname{rep}(S_1(\Lambda)).$$

Proof We know that $P_1 \cong E_1$.

We have $\nu^{-}(D(e_1\Lambda)) = \operatorname{Hom}_{\Lambda}(D(D(e_1\Lambda)), \Lambda) \cong \operatorname{Hom}_{\Lambda}(e_1\Lambda, \Lambda) \cong \Lambda e_1 = P_1$. Also

$$\nu^{-}(D(B_{1j} \otimes_{A_{j}} e_{j}\Lambda)) = \operatorname{Hom}_{\Lambda}(B_{1j} \otimes_{A_{j}} e_{j}\Lambda,\Lambda)$$

$$\cong \operatorname{Hom}_{A_{j}}(B_{1j},\operatorname{Hom}_{\Lambda}(e_{j}\Lambda,\Lambda))$$

$$\cong \operatorname{Hom}_{A_{j}}(B_{ij},\Lambda e_{j})$$

$$\cong \Lambda e_{j} \otimes_{A_{j}} \operatorname{Hom}_{A_{j}}(B_{1j},A_{j})$$

$$\cong P_{j} \otimes_{A_{j}} B_{ji}.$$
(5.1)

Since Λe_j is a finitely generated projective right A_j -module, the isomorphism in (5.1) comes from [2]. So applying the quasi-inverse Nakayama functor ν^- to (2.4), we get an exact sequence

$$0 \to \nu^{-}(E_1) \to P_1 \to \bigoplus_{j=2}^{n} P_j \otimes_{A_j} B_{j1} \to \tau^{-}(P_1) \to 0,$$
(5.2)

where $\theta_{1j} : P_1 \to P_j \bigotimes_{A_j} B_{j1}$ is given by $\lambda e_i \mapsto \sum_{r \in R_{ij}} \lambda r \otimes r^*$. We have isomorphisms $\operatorname{Hom}_{\Lambda}(P_j \bigotimes_{A_j} B_{j1}, X) \cong \operatorname{Hom}_{\Lambda}(\operatorname{Hom}_{A_j}(B_{ij}, \Lambda e_j), X) \cong B_{1j} \bigotimes_{A_j} \operatorname{Hom}_{\Lambda}(P_j, X) \cong B_{1j} \bigotimes_{A_j} X_j$. The isomorphism $\operatorname{Hom}_{\Lambda}(P_j \bigotimes_{A_j} B_{j1}, X) \to B_{1j} \bigotimes_{A_j} X_j$ is given by $f \mapsto \sum_{r \in R_{ij}} \lambda r \otimes f(e_j \otimes r^*)$, and isomorphism $\operatorname{Hom}_{\Lambda}(P_1, X) \to X_1$ is given by $g \mapsto g(e_1)$. We get a commutative diagram

$$\operatorname{Hom}_{\Lambda}(P_{j} \bigotimes_{A_{j}} B_{j1}, X) \xrightarrow{\operatorname{Hom}_{\Lambda}(\theta_{1j}, X)} \operatorname{Hom}_{\Lambda}(P_{1}, X)$$
$$\begin{array}{c} \eta_{1j}^{X} \\ & \eta_{i}^{X} \\ B_{1j} \bigotimes_{A_{j}} X_{j} \xrightarrow{\varphi_{1j}} X_{1}. \end{array}$$

This follows from that, for $f \in \operatorname{Hom}_{\Lambda}(P_j \bigotimes_{A_i} B_{j1}, X)$ and $r \in R_{1j}$, we have

$$f(r \otimes r^*) = \varphi_{1j}(r \otimes f(e_j \otimes r^*)).$$

Applying functor $\operatorname{Hom}_{\Lambda}(-, X)$ to (5.2) for $X \in \Lambda$ -mod, we obtain a commutative diagram:

Since the last two terms are isomorphic, we obtain an isomorphism $\operatorname{Hom}_{\Lambda}(\tau^{-}(P_{1}), X) \cong \operatorname{Ker}(M_{1, \operatorname{in}})$. Together with Lemma 5.3, we get the functorial isomorphism $F_{1}^{+}(-) \cong \operatorname{Hom}_{\Lambda}(T_{1}, -)$.

This theorem is the main result in this section, whose corresponding analog in [10] is [10, Theorem 9.7]. But the method for proving in [10] is incomplete for our case, the Frobenius-type triangular matrix algebra Λ .

Besides Corollary 4.13, the main results in this paper, including those in this section, are interesting to be restricted two special cases, that is, Λ is either a generalized path algebra $\Lambda = k(Q, \mathcal{A})$ endowed by Frobenius algebras A_i at each vertex $i \in Q_0$ or a path algebra $\Lambda = AQ$ of quiver Q over a Frobenius algebra A.

Acknowledgements The authors would like to thank Christof Geiss and Jan Schröer for their valuable comments and useful advices. Also, The authors warmly thank the referee for many helpful comments and suggestions in improving the quality and readability of this paper.

References

- [1] Adachi, T., Iyama, O., Reiten, I.: τ-tilting theory. Compos. Math., 150(3), 415–452 (2014)
- [2] Anderson, F. W., Fuller, K. R.: Rings and Categories of Modules, second ed., Graduate Texts in Mathematics, vol. 13, Springer-Verlag, New York, 1992
- [3] Assem, I., Simson, D., Skowroński, A.: Elements of the Representation Theory of Associative Algebras, vol.
 1, Techniques of Representation Theory, London Mathematical Society Student Texts, vol. 65, Cambridge University Press, Cambridge, 2006
- [4] Auslander, M., Platzeck, M. I., Reiten, I.: Coxeter functors without diagrams. Trans. Amer. Math. Soc., 250, 1–46 (1979)
- Bernšteřn, I. N., Gel'fand, I. M., Ponomarev, V. A.: Coxeter functors, and Gabriel's theorem. Uspehi Mat. Nauk, 282(170), 19–33 (1973)
- [6] Dlab, V., Ringel, C. M.: On algebras of finite representation type. J. Algebra, 33, 306–394 (1975)
- [7] Dlab, V., Ringel, C. M.: Indecomposable representations of graphs and algebras. Mem. Amer. Math. Soc., 6(173), v+57 (1976)
- [8] Gabriel, P.: Unzerlegbare Darstellungen. I. Manuscripta Math., 6, 71–103 (1972), correction, ibid. 6, 309 (1972)
- [9] Gabriel, P.: Auslander–Reiten sequences and representation-finite algebras. Representation theory, I (Proc. Workshop, Carleton Univ., Ottawa, Ont., 1979), Lecture Notes in Math., 831, Springer, Berlin, 1980, 1–71
- [10] Geiss, C., Leclerc, B., Schröer, J.: Quivers with relations for symmetrizable cartan matrices I : Foundations. arXiv:1410.1403, 1–67 (2015)
- [11] Iwanaga, Y.: On rings with finite self-injective dimension. Comm. Algebra, 7(4), 393-414 (1979)
- [12] Kac, V. G.: Infinite-dimensional Lie Algebras, Cambridge University Press, Cambridge, 1994
- [13] Lam, T. Y.: Serre's Problem on Projective Modules, Springer Science & Business Media, Cambridge, 2010
- [14] Li, F.: Modulation and natural valued quiver of an algebra. Pacific J. Math., 256(1), 105–128 (2012)
- [15] Li, F., Ye, C.: Gorenstein projective modules over a class of generalized matrix algebras and their applications. Algebr. Represent. Theory, 18(3), 693–710 (2015)
- [16] Li, L.: Stratifications of finite directed categories and generalized APR tilting modules. Comm. Algebra, 43(5), 1723–1741 (2015)
- [17] Nakayama, T.: On frobeniusean algebras I. Annals of Math., 611–633 (1939)
- [18] Ringel, C. M.: Representations of K-species and bimodules. J. Algebra, 41(2), 269–302 (1976)
- [19] Ringel, C. M., Zhang, P.: Representations of quivers over the algebra of dual numbers. arXiv preprint arXiv:1112.1924 (2011)
- [20] Shi, M., Li, F.: The category of modules over a triangular matrix ring of order n and its homological characterizations. Acta Math. Sin., Chin. Ser., 49(1), 215–224 (2006)
- [21] Wang, R.: Gorenstein triangular matrix rings and category algebras. J. Pure Applied Algebra, 220(2), 666–682 (2016)